


Sept London Saint Joseph Martin Saint Joseph Hospital Saint Joseph Jessamine Saint Joseph Jessamine Saint Joseph Mount Sterling Sts. Mary & Elizab ospital VNA Nazareth Home Care Flaget Memorial Hospital Frazier Rehab Institute ospital Jewish Hospital Outpatient Centers Jewish Hospital Shelbyville Jewish Physical Jewish Hospital Joseph Hospital Jeseph Hospital Jeseph Hospital Jeseph Hospital Jewish Hospital Hospital Hospital Hospital Hospital Hospital Hospital Hospital Frazier Rehab Institute Jewish Hospital Jewish Hospital Outpatie enters Jewish Hospital Shelbyville Jewish Physician Group Our Lady of Peace Saint Joseph Berea Saint Joseph Hospital Saint Joseph Jessamine Saint Joseph London Saint Joseph Martin Saint Joseph Mount Sterling Sts. Mary & Elizab ospital VNA Nazareth Home Care Flaget Memorial Hospital Frazier Rehab Institute Jewish Hospital Outpatient Centers Jewish Hospital Shelbyville Jewish Physician Group Grant Joseph Hospital Shelbyville Jewish Physician Hospital Shelbyville Jewish Physician Grant Joseph Hospital Shelbyville Jewish Physician Hospital Shelbyville Jewish Physician Grant Joseph Hospital Shelbyville Jewish Physician Grant Jewish Physician Grant Jewish Physician Grant Jewish Physician G

KentuckyOne Health Cancer Care Saint Joseph Hospital East

Public Reporting of Outcomes – 2017

2014 Cancer Program Practice Profile Reports

Saint Joseph Hospital East




BCSRT - Radiation is administered within 1 year (365 days) of diagnosis for women under the age of 70 receiving breast conservation surgery for breast cancer (Accountability)

HT - Tamoxifen or third generation aromatase inhibitor is recommended or administered within 1 year (365 days) of diagnosis for women with AJCC T1c or stage IB-III hormone receptor positive breast cancer (Accountability)

MASTRT - Radiation therapy is recommended or administered following any mastectomy within 1 year (365 days) of diagnosis of breast cancer for women with >= 4 positive regional lymph nodes (Accountability)


nBx - Image or palpation-guided needle biopsy to the primary site is performed to establish diagnosis of breast cancer (Quality Improvement)

Saint Joseph Hospital East



At least 12 regional lymph nodes are removed and pathologically examined for resected colon cancer (Quality Improvement)

Saint Joseph Hospital East



• Surgery is not the first course of treatment for cN2, M0 lung cases (Quality Improvement)